

## A Short Note on Double Roman Domination in Graphs

Abdelhak Omar<sup>1,\*</sup> and Ahmed Bouchou<sup>2</sup>

<sup>1</sup>LAMDA-RO Laboratory, Department of Mathematics, University of Blida 1, Algeria  
omar.abdelhak@etu.univ-blida.dz

<sup>2</sup>Department of Mathematics and Computer Science, University of Médéa, Algeria  
bouchou.ahmed@yahoo.fr

*Received: 14 June 2023; Accepted: 24 December 2023*

*Published Online: 11 January 2024*

**Abstract:** In this short note, we report an erroneous result of Mojdeh, Parsian and Masoumi relating the double Roman domination number to the enclaveless number and the differential of a graph.

**Keywords:** double Roman domination number, trees, differential.

**AMS Subject classification:** 05C69

### 1. Introduction

For a graph  $G = (V, E)$ , let  $\gamma(G)$ ,  $\gamma_R(G)$ ,  $\gamma_{dR}(G)$ ,  $\Psi(G)$  and  $\partial(G)$  denote the domination number, the Roman domination number, the double Roman domination number, the enclaveless number and the differential of  $G$ , respectively.

It has been shown by Mojdeh, Parsian and Masoumi [4] that for every graph  $G$  of order  $n$  having no isolated vertices,

$$\gamma_{dR}(G) \leq 2n - \Psi(G) - \partial(G) \quad (1.1)$$

It is worth noting that this result, whose invalidity will be shown, is presented in two separate papers by the same authors. The following Gallai theorems have been established in [1] and [2] for the differential of a graph and the enclaveless number, respectively.

---

\* *Corresponding Author*


**Theorem 1.** [1] If  $G$  is a graph of order  $n$ , then  $\partial(G) = n - \gamma_R(G)$ .

**Theorem 2.** [2] For any graph  $G$  of order  $n$ , then  $\Psi(G) = n - \gamma(G)$ .

Note that according to Theorems 1 and 2, the inequality (1.1) becomes  $\gamma_{dR}(G) \leq \gamma_R(G) + \gamma(G)$ . In the next section, we will provide an infinite family of graphs showing that inequality (1.1) is erroneous.

## 2. Counterexamples

Recall that a double star  $S(r, s)$  with  $r \geq s \geq 1$ , is a tree with exactly two vertices which are not leaves, one of which is adjacent to  $r$  leaves and the other one to  $s$  leaves. Let  $\mathcal{G}$  be the family of trees  $T$  obtained from a double star  $S(r, s)$  with  $r \geq s \geq 2$ , by subdividing twice the central edge and once any other edge of the double star. Figure 1 shows the smallest example of a tree belonging to  $\mathcal{G}$ . We can easily see that any tree  $T$  in  $\mathcal{G}$  has order  $n = 2(r + s) + 4$ ,  $\gamma(T) = r + s + 1$ ,  $\gamma_R(T) = r + s + 4$  and thus leading to  $\Psi(T) = r + s + 3$  and  $\partial(T) = r + s$ . Now since  $\gamma_{dR}(T) = 2(r + s) + 6$ , we consequently have  $\gamma_{dR}(T) > 2n - \Psi(T) - \partial(T)$ .




**Figure 1.** The tree  $T$  in  $\mathcal{G}$ .

In the following, we define another class of graphs different from trees for which (1.1) is not also valid. Let  $\mathcal{H}$  be the family of graphs  $G$  obtained from a star  $K_{1,p}$ , with  $p \geq 3$ , by first subdividing once each edge of the star and then adding a new vertex attached to the center vertex and one of its neighbors. Figure 2 shows the smallest example of a graph belonging to  $\mathcal{H}$ . One can easily see that any graph  $G$  in  $\mathcal{H}$  has order  $n = 2p + 2$ ,  $\gamma(G) = p$ ,  $\gamma_R(G) = p + 2$  and thus leading to  $\Psi(G) = p + 2$  and  $\partial(G) = p$ . Now since  $\gamma_{dR}(G) = 2p + 3$ , we consequently have  $\gamma_{dR}(G) > 2n - \Psi(G) - \partial(G)$ .

We conclude by mentioning that inequality (1.1) is used in [3], which therefore calls into question the validity of certain results.

**Acknowledgements.** The authors are thankful to the anonymous reviewers for their valuable comments and suggestions.



**Figure 2.** The graph  $G$  in  $\mathcal{H}$ .

**Conflict of interest.** The authors declare that they have no conflict of interest.

**Data Availability.** Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

## References

- [1] S. Bermudo, H. Fernau, and J. Sigarreta, *The differential and the Roman domination number of a graph*, Appl. Anal. Discret. Math. **8** (2014), no. 1, 155–171.
- [2] J.R. Lewis, *Differentials of Graphs*, East Tennessee State University, Johnson City, 2004.
- [3] D.A. Mojdeh, I. Masoumi, and A. Parsian, *A new approach on Roman graphs*, Turk. J. Math. Comput. Sci. **13** (2021), no. 1, 6–13.
- [4] D.A. Mojdeh, A. Parsian, and I. Masoumi, *Characterization of double Roman trees*, Ars Combin. **153** (2020), 53–68.