

## Two techniques to reduce the Pareto optimal solutions in multi-objective optimization problems

Fatemeh Ahmadi<sup>†</sup>, Davoud Foroutannia<sup>\*</sup>

Department of Mathematics, Vali-e-Asr University of Rafsanjan, Rafsanjan, Iran

<sup>†</sup>[fatemeh.Ahmadi@stu.vru.ac.ir](mailto:fatemeh.Ahmadi@stu.vru.ac.ir)

<sup>\*</sup>[foroutan@vru.ac.ir](mailto:foroutan@vru.ac.ir)

*Received: 15 June 2023; Accepted: 8 May 2024*

*Published Online: 7 June 2024*

**Abstract:** In this study, for a decomposed multi-objective optimization problem, we propose the direct sum of the preference matrices of the subproblems provided by the decision maker (DM). Then, using this matrix, we present a new generalization of the rational efficiency concept for solving the multi-objective optimization problem (MOP). A problem that sometimes occurs in multi-objective optimization is the existence of a large set of Pareto optimal solutions. Hence, decision making based on selecting a unique preferred solution becomes difficult. Considering models with the concept of generalized rational efficiency can relieve some of the burden from the DM by shrinking the solution set. This paper discusses both theoretical and practical aspects of rationally efficient solutions related to this concept. Moreover, we present two techniques to reduce the Pareto optimal solutions using. The first technique involves using the powers of the preference matrix, while the second technique involves creating a new preference matrix by modifying the decomposition of the MOP.

**Keywords:** pareto, nondominated,  $A_P$ -efficiency, multi-objective programming

**AMS Subject classification:** 90C29, 91B08, 90B50.

### 1. Introduction

Because of the conflicting nature of the objective functions, a MOP usually does not have a single optimal solution for all objectives, but a set of Pareto optimal solutions exists. Several multi-objective optimization approaches exist that generate subsets of Pareto optimal solutions, which can be overwhelming to the DM in the task of selecting the most appropriate solution to implement. More recently, these approaches have categorized into a priori, a posteriori, interactive, and pruning methods by Petchrompo et al. [12].

---

<sup>\*</sup> *Corresponding Author*

In the a priori method, a MOP is converted into a single objective problem before obtaining a solution. A priori methods use DM preferences to bias the search of optimal solutions towards a preferred region, for example by changing the definition of dominance [7, 9, 19], by giving different weights to the objectives [17], by assigning reference values (goals) and priority levels to the objectives [18], by assuming a utility function describing the DM behaviour and interest in the alternative solutions [8]. The a posteriori method generates a representative set of Pareto optimal solutions and the DM chooses the best one among them, [16, 20]. Interactive methods allow the DM to guide the search by alternating optimization, and preference articulation iteratively, [8, 15]. Pruning methods are applied before, during, or after the optimization process to reduce the number of Pareto optimal solutions, [12, 13].

One approach to reduce the size of the Pareto optimal set is to combine multiple objectives, i.e. by summing them, before employing tools to solve the resulting MOP, [2]. This method also reduces the objective space dimension using DM's preference matrix. In particular, it can be used to discard certain unwanted solutions, especially the extreme points which are found by minimizing just one of the objectives given in the classical sense while disregarding the rest. Unfortunately, the problem of combining several objectives does not seem to have attracted too much attention in the literature so far.

Berman and Naumov [1] were perhaps the first to construct a matrix of a cone to represent DM's preferences by interval trade-offs. Noghin [10, 11] investigated the relative importance of objectives and provided a definition of weights (which he called coefficients) for objective functions as well as for groups of objective functions. Podinowski [14] extended the concept of "one criterion is more important than another" to "one criterion is n-times more important than another" by applying pairwise coefficients of relative importance obtained from the DM. Also, using these coefficients, he constructed a matrix that represents DM's preferences. The polyhedral cone and the partial order characterized by it are discussed by Engau and Wiecek [3]. They discussed the relationship between the optimal solutions with respect to a polyhedral cone represented by a matrix  $A$  and the optimal solutions of a multi-objective optimization with respect to the natural order after linear transformation of the objective functions by a matrix  $A$ . Hunt et al. [5] developed a convex polyhedral cone-based preference modeling framework for decision making with multiple criteria, which extends the classical notion of Pareto optimality and accounts for the relative importance of criteria. Dempe et al. [2] investigated the impact of using linear combinations of objectives and which solutions are eliminated by doing so. They showed how the strategy of combining objectives linearly influences the efficient set.

In this paper, we do not focus on constructing a preference matrix. Instead, we investigate two pruning techniques to reduce the number of Pareto optimal solutions, by applying the strategy of linear combination of objectives. The present study is an extension of some results obtained by [6, 7].

The paper is organized as follows. In Section 2, we give some basic and preliminary concepts. In Section 3, we introduce the concept of rational  $A_P$ -efficiency, and we define rational  $A_P$ -efficiency in terms of vector inequalities, in order to make it prac-

tical. In Section 4, the concept of  $A_P^r$ -efficiency is examined to generate a subset of Pareto optimal solutions, for  $r = 1, 2, \dots$ . Moreover, the  $A_P$ -efficient solutions are reduced by constructing a new preference matrix which based on  $P$  and  $A_P$ , in Section 5. Also, an algorithm is presented to generate a subset of  $A_P$ -efficient solutions. In addition, the numerical examples are provided to confirm the efficiency of these methods. Finally, Section 6 concludes the paper.

## 2. Terminology

In this article, the following notations will be used. Let  $\mathbb{R}^m$  be the Euclidean vector space and  $y', y'' \in \mathbb{R}^m$ .  $y' \preceq y''$  denotes  $y'_i \leq y''_i$  for all  $i = 1, \dots, m$ .  $y' < y''$  denotes  $y'_i < y''_i$  for all  $i = 1, \dots, m$ .  $y' \preceq y''$  denotes  $y' \preceq y''$  but  $y' \neq y''$ .

Consider a decision problem defined as an optimization problem with  $m$  objective functions. For simplification, we assume, without losing generality, that the objective functions should be minimized. The problem can be defined as follows:

$$\begin{aligned} & \min (f_1(x), \dots, f_m(x)) \\ & \text{subject to } x \in X, \end{aligned} \tag{2.1}$$

where  $x$  denotes a vector of decision variables selected from the feasible set  $X$  and  $f(x) = (f_1(x), \dots, f_m(x))$  is a vector function that maps the feasible set  $X$  into the objective (criterion) space  $\mathbb{R}^m$ . We refer to the elements of the objective space as outcome vectors. An outcome vector  $y$  is attainable if it expresses the outcomes of a feasible solution, i.e.,  $y = f(x)$  for some  $x \in X$ . The set of all attainable outcome vectors will be denoted by  $Y = f(X)$ .

In single objective minimization problems, we compare the objective values at different feasible decisions to select the best decision. Decisions are ranked according to the objective values at those decisions and the decision resulting in the least smallest objective value is the most preferred decision. Similarly, to make the multi-objective optimization model operational, one needs to assume some solution concept specifying what it means to minimize multi-objective functions. The solution concepts are defined by the properties of the corresponding preference model. We assume that solution concepts depend only on the evaluation of the outcome vectors while not taking into account any other solution properties not represented within the outcome vectors. Thus, we can limit our considerations to the preference model in the objective space  $Y$ . In the following, some basic concepts and definitions of preference relations are reviewed from [6].

**Definition 1.** Let  $y', y'' \in \mathbb{R}^m$  and let  $\preceq$  be a relation of weak preference defined on  $\mathbb{R}^m \times \mathbb{R}^m$ . The corresponding relations of strict preference  $\prec$  and indifference  $\simeq$  are defined as follows:

$$y' \prec y'' \Leftrightarrow (y' \preceq y'' \text{ and not } y'' \preceq y'), \tag{2.2}$$

$$y' \simeq y'' \Leftrightarrow (y' \preceq y'' \text{ and } y'' \preceq y'). \tag{2.3}$$

**Definition 2.** Preference relations satisfying the following axioms are called rational preference relations:

1. Reflexivity: for all  $y \in \mathbb{R}^m$ :  $y \preceq y$ .
2. Transitivity: for all  $y', y'', y''' \in \mathbb{R}^m$ :  $y' \preceq y''$  and  $y'' \preceq y''' \Rightarrow y' \preceq y'''$ .
3. Strict monotonicity: for all  $y \in \mathbb{R}^m$ :  $y - \epsilon e_i \prec y$  for  $\epsilon > 0$  where  $e_i$  denotes the  $i^{\text{th}}$  unit vector in  $\mathbb{R}^m$ .

The rational preference relations allow us to formalize the Pareto optimal solution concept with the following definitions.

**Definition 3.** Let  $y', y'' \in Y$ . We say that  $y'$  rationally dominates  $y''$ , and denote by  $y' \prec_r y''$  iff  $y' \prec y''$  for all rational preference relations  $\preceq$ . An outcome vector  $y$  is rationally nondominated if and only if there does not exist another outcome vector  $y'$  such that  $y' \prec_r y$ . Analogously, a feasible solution  $x \in X$  is a rationally efficient solution of the MOP (2.1) if and only if  $y = f(x)$  is rationally nondominated.

A relationship between the weak rational preference relation  $\preceq_r$  and the Pareto relation has been established in [6]. As the following proposition shows finding nondominated points with respect to the relation  $\preceq_r$  can be done by means of Pareto preference.

**Proposition 1.** ([6], Proposition 1.1) For any two vectors  $y', y'' \in Y$ , we have

$$\begin{aligned} y' \preceq_r y'' &\Leftrightarrow y' \leq y'', \\ y' \prec_r y'' &\Leftrightarrow y' \leq y''. \end{aligned}$$

The set of all rationally efficient solutions  $x \in X$  is denoted by  $X_E$  and called the efficient set. The set of all rationally nondominated points  $y = f(x) \in Y$ , where  $x \in X_E$ , is denoted by  $Y_N$  and called the nondominated set.

### 3. The concept of rational $A_P$ -efficiency

In this section, we will introduce a dominance relation to generate solutions that are rationally  $A_P$ -efficient. The following definitions are necessary for the concepts of interest in this paper.

**Definition 4.** Let  $M = \{1, \dots, m\}$  be the index set of objective functions  $f = (f_1, \dots, f_m)$  and  $n$  be a positive integer such that  $n \leq m$ . A collection  $P = \{P_k \subseteq M : k = 1, \dots, n\}$  is called a decomposition of  $M$ , and also it is said a partition of  $M$  if  $\bigcup_{k=1}^n P_k = M$ , and  $P_i \cap P_j = \emptyset$  for all  $i \neq j$ , where  $i, j \in \{1, \dots, n\}$  and  $P_k$  is index set of objective functions in class  $k$ . The multi-objective problem

$$\begin{aligned} \min(f_j(x))_{j \in P_k} \quad & (k = 1, \dots, n), \\ \text{subject to } x \in X, \end{aligned} \tag{3.1}$$

is called a subproblem of the multi-objective problem (2.1) and the collection of all these subproblems is called a decomposition of the multi-objective problem (2.1).

To simplify the notations, we put  $f^k = (f_j)_{j \in P_k}$  and  $f^k(x) = (f_j(x))_{j \in P_k}$ . Also, we will use the notation  $y = (y_{P_k})_{k=1}^n$ , where  $y_{P_k} = (y_i)_{i \in P_k}$ , for any vector  $y = (y_1, \dots, y_m) \in \mathbb{R}^m$ . It is noteworthy that the functions  $f^k$  maps the feasible set  $X$  into the space  $\mathbb{R}^{|P_k|}$ , where  $|P_k|$  denotes the cardinality of the set  $P_k$  for  $k = 1, \dots, n$ , and  $m = |P_1| + \dots + |P_n|$ .

The preference information identifies the relative importance of different objectives in the optimization problem. The DM can select the preference matrix  $A_k$  that displays the relative importance of different objectives in the subproblem (3.1) as follows. Let  $i, j \in P_k$ ,  $i \neq j$  and criterion  $j$  be relatively more important than criterion  $i$ , so that an improvement of the former is desired even when the latter decays. Let  $a_{ij}^k \geq 0$  represent the minimum desired improvement in the  $j$ -th objective function for a unit loss in the  $i$ -th objective function. If criteria  $i$  and  $j$  are equally important than there is no specific preference relation between them and the corresponding coefficient  $a_{ij}^k = 0$ . For  $i = j$ , let  $a_{ij}^k = 1$ . The elements  $a_{ij}^k$  form a  $|P_k| \times |P_k|$  matrix  $A_k = [a_{ij}^k]$ . For the case of Pareto domination, the matrix  $A_k$  is an identity matrix with  $a_{ij}^k = 1$  for  $i = j$  and 0 otherwise. In general, the preference matrix represents a polyhedral cone that plays the role of a domination cone for the used preference model, which is based on the trade-off between objectives. For more details, the reader can refer to [1–3, 5]. In this paper, we do not focus on how to construct a preference matrix that is provided by a DM. Instead, we suggest two pruning methods that are based on preference matrix. The idea behind these is that the DM classifies the objective functions in different classes and determines a partition  $P = \{P_k \subseteq M : k = 1, \dots, n\}$  of  $\{1, \dots, m\}$  according to the importance of objective functions. The DM should provide the preference matrix  $A_k$  for the relative importance of the objectives of subproblem (3.1) for  $k = 1, \dots, n$ . Thus, we introduce the matrix  $A_P = A_1 \oplus \dots \oplus A_n$ , which is the direct sum of the matrices  $A_1, \dots, A_n$ , i.e.

$$A_P = \begin{bmatrix} A_1 & 0 & \dots & 0 \\ 0 & A_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & A_n \end{bmatrix}.$$

We have  $A_P(y) = (A_1(y_{P_1}), \dots, A_n(y_{P_n}))$  for  $y \in Y$ , where  $y_{P_k} = (y_j)_{j \in P_k}$  for  $k = 1, \dots, n$ .

**Definition 5.** Let  $y', y'' \in Y$  be two outcome vectors. We say that  $y'$  rationally  $A_P$ -dominates  $y''$ , and denote by  $y' \prec_{rA_P} y''$  if and only if  $A_P(y') \prec A_P(y'')$  for all rational preference relations  $\preceq$ . An outcome vector  $y$  is called rationally  $A_P$ -nondominated if and only if there is not another outcome vector  $y'$  such that  $y' \prec_{rA_P} y$ . Analogously, a feasible solution  $x \in X$  is called a rationally  $A_P$ -efficient solution of the problem (2.1) if and only if  $y = f(x)$  is rationally  $A_P$ -nondominated point.

The set of all rationally  $A_P$ -efficient solutions  $x \in X$  is denoted by  $X_{A_P E}$  and is called the rationally  $A_P$ -efficient set. The set of all rationally  $A_P$ -nondominated points  $y = f(x) \in Y$ , where  $x \in X_{A_P E}$ , is denoted by  $Y_{A_P N}$  and is called the rationally  $A_P$ -nondominated set.

Let  $A$  be the preference matrix representing the relative importance of different objectives in the problem (2.1). It should be noted that  $A_P = A$ , when  $n = 1$  or  $P_1 = \{1, \dots, m\}$ . Therefore, we can rewrite the above definition for the matrix  $A$  as follows.

**Definition 6.** Let  $y', y'' \in Y$ . We say that  $y'$  rationally  $A$ -dominates  $y''$ , and denote by  $y' \prec_{rA} y''$  if and only if  $A(y') \prec A(y'')$  for all rational preference relations  $\preceq$ . An outcome vector  $y$  is called rationally  $A$ -nondominated if and only if there is not another outcome vector  $y'$  such that  $y' \prec_{rA} y$ . Analogously, a feasible solution  $x \in X$  is called a rationally  $A$ -efficient solution of the problem (2.1) if and only if  $y = f(x)$  is rationally  $A$ -nondominated point.

It is worth noting that for  $n = 1$  or  $P_1 = \{1, \dots, m\}$ , we have  $X_{A_P E} = X_{AE}$  and  $Y_{A_P N} = Y_{APN}$ . Let  $I$  denote the identity matrix. If  $P_k = \{k\}$  and  $A_k = I$  for all  $k = 1, \dots, m$ , then the relation  $\prec_{rA_P}$  becomes the rational relation  $\prec_r$ .

Similar to the relation  $\prec_{rA_P}$ , we can define the relation of rational  $A_P$ -indifference,  $\simeq_{rA_P}$ , and the relation of rational weak  $A_P$ -dominance,  $\preceq_{rA_P}$ . We say that  $y' \simeq_{rA_P} y''$  if and only  $A_P(y') \simeq A_P(y'')$  for all rational preference relations  $\preceq$ , and also  $y' \preceq_{rA_P} y''$  if and only  $A_P(y') \preceq A_P(y'')$  for all rational preference relations  $\preceq$ . The relations  $\prec_{rA_P}$ ,  $\simeq_{rA_P}$  and  $\preceq_{rA_P}$  satisfy conditions (2.2-2.3).

It is clear that the preference relation  $\preceq_{rA_P}$  satisfies the reflexivity and transitivity axioms. To continue, we express some conditions that guarantee the relation  $\preceq_{rA_P}$  is a rational preference relation. Throughout this section, we assume that  $e_i^k \in \mathbb{R}^k$  is the unit vector with the  $i$ -th component equal to one and the remaining ones equal to zero, where  $k = 1, \dots$  and  $i \in \{1, \dots, k\}$ .

**Theorem 1.** *The strict monotonicity axiom for the preference  $\preceq_{rA_P}$  is equivalent to the condition*

$$a_i^k \geq 0 \quad (i = 1, \dots, |P_k|), \quad (3.2)$$

where  $a_i^k$  is the column  $i$  of the matrix  $A_k$  for  $k = 1, \dots, n$ .

*Proof.* We will prove that if matrix  $A_p$  satisfies condition (3.2), then the strict monotonicity axiom holds for preference  $\preceq_{rA_P}$ . For  $y \in Y$ ,  $i \in \{1, \dots, m\}$ ,  $\epsilon > 0$  and  $y' = y - \epsilon e_i^m$ , we show that  $y' \prec_{rA_P} y$ . There exists an index  $k \in \{1, \dots, n\}$  such that  $i \in P_k$ . It's obvious that  $y'_{P_j} = y_{P_j}$  and  $A_j(y'_{P_j}) = A_j(y_{P_j})$ , when  $j \in \{1, \dots, n\} - \{k\}$ .

Since  $y'_{P_k} = y_{P_k} - \epsilon e_i^{|P_k|}$ , we have  $y'_{P_k} \leq y_{P_k}$ . According to condition (3.2), we obtain

$$\sum_{j=1}^{|P_k|} a_{ij}^k (y'_{P_k})_j \leq \sum_{j=1}^{|P_k|} a_{ij}^k (y_{P_k})_j \quad (i = 1, \dots, |P_k|),$$

where strict inequality holds at least once. So  $A_k(y'_{P_k}) \leq A_k(y_{P_k})$  and the proof is complete by applying Proposition 1.

Conversely, assuming that the preference  $\preceq_{rA_P}$  follows the strict monotonicity axiom. We have

$$e_j^m - e_j^m \prec_{eA_P} e_j^m, \quad (j = 1, \dots, m),$$

which, together with Proposition 1, imply that matrix  $A_P$  satisfies condition (3.2).  $\square$

To make it practical, rational  $A_P$ -efficiency is defined in terms of vector inequalities. To do that, we define a certain preference relation.

**Definition 7.** Suppose that  $y', y'' \in Y$  are two outcome vectors. We define the relation  $\leq_{A_P}$  as follows:

$$y' \leq_{A_P} y'' \Leftrightarrow A_P(y') \leq A_P(y''). \quad (3.3)$$

Also, we can define the relations  $<_{A_P}$  and  $=_{A_P}$  as follows:

$$\begin{aligned} y' <_{A_P} y'' &\Leftrightarrow (y' \leq_{A_P} y'' \text{ and not } y'' \leq_{A_P} y'), \\ y' =_{A_P} y'' &\Leftrightarrow (y' \leq_{A_P} y'' \text{ and } y'' \leq_{A_P} y'). \end{aligned}$$

It is clear that the preference relation  $\leq_{A_P}$ , satisfies reflexivity, transitivity and monotonicity. This means that, the relation (3.3) is a rational preference relation. Note that the relation  $\leq_{A_P}$  becomes the Pareto relation, when  $A = I$  and  $P_k = \{k\}$  for all  $k = 1, \dots, m$ .

In the following, we will discuss the relationship between two preferences  $\preceq_{rA_P}$  and  $\leq_{A_P}$ .

**Corollary 1.** Let  $y', y'' \in Y$  be two outcome vectors. We have

$$y' \preceq_{rA_P} y'' \Leftrightarrow y' \leq_{A_P} y'',$$

$$y' \prec_{rA_P} y'' \Leftrightarrow y' <_{A_P} y''.$$

*Proof.* Based on Definition 5, Proposition 1 and Definition 7, we can conclude that

$$\begin{aligned} y' \preceq_{r_{A_P}} y'' &\Leftrightarrow A_P(y') \preceq_r A_P(y'') \\ &\Leftrightarrow A_P(y') \leq A_P(y'') \\ &\Leftrightarrow y' \leq_{A_P} y''. \end{aligned}$$

□

The statement below is true according to Theorem 1 and Corollary 1.

**Corollary 2.** *The preference  $\leq_{A_P}$  is a rational preference relation if and only if the matrix  $A_P$  satisfies condition (3.2).*

Note that Corollary 1 permits one to express rational  $A_P$ -efficiency for problem (2.1) in terms of the standard efficiency for the MOP with objectives  $A_P(f(x))$ ,

$$\begin{aligned} &\min A_P(f(x)) \\ &\text{subject to } x \in X. \end{aligned} \tag{3.4}$$

**Corollary 3.** *A feasible solution  $x \in X$  is a rationally  $A_P$ -efficient solution to the problem (2.1) if and only if it is an efficient solution to the problem (3.4).*

**Remark 1.** If  $P_1 = \{1, \dots, m\}$  in Corollary 3, then any feasible solution  $x \in X$  is a rationally  $A$ -efficient solution of the problem (2.1), if and only if it is an efficient solution of the problem

$$\begin{aligned} &\min A(f(x)) \\ &\text{subject to } x \in X. \end{aligned} \tag{3.5}$$

Also, if  $A_k = I$  is identity matrix for all  $k = 1, \dots, n$ , then we have Corollary 3.2 from [7]. In addition, if  $A_k = I$  and  $P_k = \{k\}$  for all  $k = 1, \dots, m$ , we have Proposition 1.1 from [6].

**Example 1.** Let

$$X = \{(x_1, x_2, x_3) \in \mathbb{R}^3 : x_1 + x_2 \geq 1 \text{ and } x_1, x_2, x_3 \geq 0\},$$

and  $f(x_1, x_2, x_3) = (x_1, x_2, x_3)$  and  $P_1 = \{1, 2\}$  and  $P_2 = \{3\}$ . We get

$$X_E = \{(x_1, x_2, x_3) \in \mathbb{R}^3 : x_1 + x_2 = 1 \text{ and } x_1, x_2 \geq 0, x_3 = 0\}.$$

For  $A_1 = \begin{bmatrix} 1 & 0 \\ \alpha & 1 \end{bmatrix}$  and  $A_2 = [1]$ , we have  $A_P(f(x)) = (x_1, \alpha x_1 + x_2, x_3)$ , where  $\alpha \geq 0$  and  $x \in X$ . The assumption  $x_1 + x_2 = 1$  implies that  $\alpha x_1 + x_2 = (\alpha - 1)x_1 + 1$ . Hence, it is easy to obtain that  $X_{A_P E} = X_E$ , when  $0 \leq \alpha < 1$ .

Now, let  $\alpha \geq 1$  and  $\hat{x} \in X$  be a feasible solution with  $\hat{x}_1 > 0$ . For  $0 < \epsilon < \hat{x}_1$ , we put  $x_1 = \hat{x}_1 - \epsilon$ ,  $x_2 = \hat{x}_2 + \epsilon$  and  $x_3 = \hat{x}_3$ . After some calculations, we conclude that  $A_P(f(x)) \leq A_P(f(\hat{x}))$ , and thus  $X_{A_P E} = \{(0, 1, 0)\}$ .



Finally, we show that rational  $A_P$ -efficiency is scale invariant with respect to linear scaling with a positive factor.

**Theorem 2.** *Let  $g : R \rightarrow R$  be a strictly increasing linear function. The feasible solution  $x \in X$  is a rationally  $A_P$ -efficient solution of the problem (2.1), if and only if it is a rationally  $A_P$ -efficient solution of the problem*

$$\min_{x \in X} (g(f_1(x)), \dots, g(f_m(x))). \quad (3.6)$$

*Proof.* Let  $x$  be a rationally  $A_P$ -efficient solution of the problem (2.1). If  $x$  is not a rationally  $A_P$ -efficient solution of the problem (3.6). Then there is a feasible vector  $x'$  such that the outcome vectors  $y = (g(f_1(x)), \dots, g(f_m(x)))$  and  $y' = (g(f_1(x')), \dots, g(f_m(x')))$  satisfy

$$A_k(y'_{P_k}) \leq A_k(y_{P_k}) \quad (k = 1, \dots, n),$$

and  $A_k(y'_{P_k}) \leq A_k(y_{P_k})$  for some  $k \in \{1, \dots, n\}$ . Since the function  $g$  is strictly increasing and linear, we obtain

$$A_k(f_{P_k}(x')) \leq A_k(f_{P_k}(x)) \quad (k = 1, \dots, n),$$

and  $A_k(f_{P_k}(x')) \leq A_k(f_{P_k}(x))$  for some  $k \in \{1, \dots, n\}$ . Hence  $f(x')$  rationally  $A_P$ -dominates  $f(x)$ , which contradicts the rationally  $A_P$ -efficient of  $x$ . The proof of converse part can be done in a similar way.  $\square$

#### 4. The concept of $A_P^r$ -efficiency

In this section, we investigate the relationship between  $A_P$ -efficient solutions and  $A_P^r$ -efficient solutions. Then, we introduce the concept of  $A_P^\infty$ -efficient to generate a subset of efficient solutions, which aims to offer a limited number of representative solutions to the DM. At first, we study the relationship between  $A_P$ -efficient solutions and  $(AB)_P$ -efficient solutions. To do this, we require the following theorem.

**Theorem 3.** *Let  $A = (a_1, \dots, a_m)$  and  $B = (b_1, \dots, b_m)$  be two  $m \times m$  matrices, where  $a_j$  and  $b_j$  are column  $j$  of the matrices  $A$  and  $B$ , respectively.*

(i) *Let  $AB = C = (c_1, \dots, c_m)$ , where  $c_j$  is column  $j$  of the matrix  $C$ . If  $a_j \geq 0$  and  $b_j \geq 0$  for all  $j = 1, \dots, m$ , then  $c_j \geq 0$  for all  $j = 1, \dots, m$ .*

(ii) *If  $a_j \geq 0$  for all  $j = 1, \dots, m$  and  $y', y''$  are two outcome vectors in  $R^m$ , then*

$$\begin{aligned} y' <_B y'' &\implies y' <_{AB} y'', \\ y' \leq_B y'' &\implies y' \leq_{AB} y''. \end{aligned}$$

*Hence  $Y_{(AB)N} \subset Y_{BN}$ . This implies that  $X_{(AB)E} \subset X_{BE}$ . In particular, we have  $X_{AE} \subset X_E$ , when  $B = I$ .*

*Proof.* (i) We have

$$c_j = \left( \sum_{k=1}^m a_{ik} b_{kj} \right)_{i=1}^m.$$

The condition  $b_j \geq 0$  implies that  $b_{kj} \geq 0$  for all  $k = 1, \dots, m$  and  $b_{k'j} > 0$  for some  $k' \in \{1, \dots, m\}$ . Also  $a_{k'} \geq 0$  conclude that  $a_{ik'} \geq 0$  for any  $i = 1, \dots, m$  and  $a_{i'k'} > 0$  for some  $i' \in \{1, \dots, m\}$ . Thus  $a_{ik} b_{kj} \geq 0$  for any  $i, k = 1, \dots, m$  and  $a_{i'k'} b_{k'j} > 0$ , which means that  $c_j \geq 0$ .

(ii) Let  $y', y'' \in R_{\geq}^m$  and  $y' <_B y''$ , hence

$$\sum_{j=1}^m b_{kj} y'_j \leq \sum_{j=1}^m b_{kj} y''_j \quad (k = 1, \dots, m),$$

and  $\sum_{j=1}^m b_{k'j} y'_j < \sum_{j=1}^m b_{k'j} y''_j$  for some  $k' \in \{1, \dots, m\}$ . Now according to assumption, we have  $a_{ik} \geq 0$  for all  $i, k = 1, \dots, m$ , and also there exists  $i' \in \{1, \dots, m\}$  such that  $a_{i'k'} > 0$ . This implies that

$$\begin{aligned} \sum_{k=1}^m a_{ik} \sum_{j=1}^m b_{kj} y'_j &\leq \sum_{k=1}^m a_{ik} \sum_{j=1}^m b_{kj} y''_j \quad (i = 1, \dots, m), \\ \Rightarrow \sum_{j=1}^m \left( \sum_{k=1}^m a_{ik} b_{kj} \right) y'_j &\leq \sum_{j=1}^m \left( \sum_{k=1}^m a_{ik} b_{kj} \right) y''_j \quad (i = 1, \dots, m), \end{aligned}$$

where the inequality is strict for  $i = i'$ . Thus  $AB y' \leq AB y''$ , which means that  $y' <_{AB} y''$ . □

**Theorem 4.** Let  $A_P = A_1 \oplus \dots \oplus A_n$  and  $B_P = B_1 \oplus \dots \oplus B_n$  be two  $m \times m$  matrices and  $a_j \geq 0$  for  $j = 1, \dots, m$ , where  $a_j$  is column  $j$  of the matrix  $A_P$ . If  $y', y''$  are two outcome vectors in  $R^m$  with  $y' <_{B_P} y''$ , then  $y' <_{(AB)_P} y''$ , hence  $Y_{(AB)_P N} \subset Y_{B_P N}$ . This concludes that  $X_{(AB)_P E} \subset X_{B_P E}$ .

*Proof.* Let  $y', y'' \in Y$  and  $y' <_{B_P} y''$ . Hence

$$B_k(y'_{P_k}) \leq B_k(y''_{P_k}) \quad (k = 1, \dots, n),$$

and  $B_k(y'_{P_k}) < B_k(y''_{P_k})$  for some  $k \in \{1, \dots, n\}$ . Let  $a_j^k$  be column  $j$  of the matrix  $A_k$  for  $k = 1, \dots, n$ . Since  $a_j^k \geq 0$ , by using Theorem 3, we have

$$A_k B_k(y'_{P_k}) \leq A_k B_k(y''_{P_k}) \quad (k = 1, \dots, n),$$

and  $A_k B_k(y'_{P_k}) < A_k B_k(y''_{P_k})$  for some  $k \in \{1, \dots, n\}$ . This implies that  $y' <_{(AB)_P} y''$ . □

The following statement states the relationship among  $A_P$ -efficient solutions,  $A$ -efficient solutions and efficient solution of problem (2.1).

**Corollary 4.** *Let  $A_P = A_1 \oplus \dots \oplus A_n$  be an  $m \times m$  matrix and  $a_j \geq 0$  for  $j = 1, \dots, m$ , where  $a_j$  is column  $j$  of the matrix  $A_P$ . If  $y', y'' \in Y$  and  $y' \leq y''$ , then  $y' <_{A_P} y''$ , hence  $Y_{A_P N} \subset Y_N$ . This concludes that  $X_{A_P E} \subset X_E$ . Thus if  $x \in X$  is an efficient solution of problem (3.4), then it is an efficient solution of problem (2.1). In particular, If  $n = 1$ , then  $Y_{AN} \subset Y_N$  and  $X_{AE} \subset X_E$ . Thus, if  $x \in X$  is an efficient solution of problem (3.5), then it is an efficient solution of problem (2.1).*

*Proof.* Setting  $B_P = I$  in the above theorem gives the desired result.  $\square$

For a square matrix  $A$  and a positive integer  $r$ ,  $A^r$  represents the product of  $A$  multiplied by itself  $r$  times. We also define  $A^0 = I$ .

**Corollary 5.** *Suppose that the matrix  $A_P$  satisfies the condition  $a_j \geq 0$  for  $j = 1, \dots, m$ . If  $r$  is a non-negative integer, then  $Y_{A_P^{r+1} N} \subset Y_{A_P^r N} \subset Y_N$ .*

*Proof.* The result follows by replacing  $A^r$  instead of  $B$ , in Theorem 4.  $\square$

Let  $\mathbb{R}_{\geq}^m = \{d \in \mathbb{R}^m : d \geq 0\}$  be non-negative orthant of  $\mathbb{R}^m$ . By Corollary 3, one can easily verify that the feasible solution  $\hat{x} \in X$  is a rationally  $A_P^r$ -efficient solution if and only if

$$(A_P^r(f(\hat{x})) - \mathbb{R}_{\geq}^m) \cap A_P^r(f(X)) = \{A_P^r(f(\hat{x}))\}. \quad (4.1)$$

The condition  $a_j \geq 0$  for  $j = 1, \dots, m$ , in the above results are necessary. To investigate this fact, we give the following example.

**Example 2.** Let

$$X = Y = \{(y_1, y_2) : y_1^2 + y_2^2 \leq 1\},$$


and  $A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$  and  $P_1 = \{1, 2\}$ . Using the relation (4.1) for  $r = 0$  and  $r = 1$ , we have

$$Y_N = \{(y_1, y_2) : y_1^2 + y_2^2 = 1, -1 \leq y_1, y_2 \leq 0\},$$

and

$$Y_{AN} = \{(y_1, y_2) : y_1^2 + y_2^2 = 1, -1 \leq y_1 \leq 0, 0 \leq y_2 \leq 1\},$$

respectively. Since  $A^{2r} = I$  and  $A^{2r+1} = A$  for  $r = 0, 1, \dots$ , we get  $Y_{A^{2r} N} = Y_N$  and  $Y_{A^{2r+1} N} = Y_{AN}$ .



**Figure 1.** The  $A_P^r$ -efficient solutions of the test problem (4.2) for  $r = 0, 1, 2, 7$ .

In the following example, we investigate Corollary 5, and show that the set of  $A_P^r$ -efficient solutions are reducing when  $r$  is increasing. For this purpose, a large number of random solutions are generated for scalable test function. From this large set of solutions, efficient solutions and  $A_P^r$ -efficient are calculated for  $r = 1, 2, 7$ .

**Example 3.** Let us consider the following test problem from [4],

$$\begin{aligned}
 \min_{x \in \mathbb{R}^2} \quad & y = (f_1(x), f_2(x), f_3(x), f_4(x), f_5(x), f_6(x)) \\
 & f_1(x) = x_1^2 + (x_2 + 1)^2 \\
 & f_2(x) = (x_1 - 0.5)^2 + (x_2 + 0.5)^2 \\
 & f_3(x) = (x_1 - 1)^2 + x_2^2 \\
 & f_4(x) = (x_1 + 1)^2 + x_2^2 \\
 & f_5(x) = (x_1 - 0.5)^2 + (x_2 - 0.5)^2 \\
 & f_6(x) = x_1^2 + (x_2 - 1)^2 \\
 & x_1, x_2 \in [-1, 1].
 \end{aligned} \tag{4.2}$$

To solve this problem, we use MATLAB software environment. We select 3000 random solutions  $(x_1, x_2) \in [-1, 1] \times [-1, 1]$ , and compare the values of the objective functions at these solutions. Among these solutions, there are 1741 efficient solutions, which are indicated by the blue points in Figure 1. Let  $P_1 = \{1, 2, 3\}$ ,  $P_2 = \{4, 5, 6\}$  and

$$A_1 = \begin{bmatrix} 1 & 0.5 & 0 \\ 0.9 & 0.8 & 0.5 \\ 1 & 0 & 0 \end{bmatrix}, \quad A_2 = \begin{bmatrix} 1 & 0.7 & 0 \\ 1 & 0 & 0.8 \\ 0 & 0.5 & 0.8 \end{bmatrix},$$

be given by the DM. By comparing the values of  $A_P^r(f(x))$  at the efficient solutions for  $r = 1, 2, 7$ , we obtain 888  $A_P$ -efficient solutions, 420  $A_P^2$ -efficient solutions and 129  $A_P^7$ -efficient solutions, which are shown by red circles, green square and cyan star, respectively, in Figure 1.

Using the results above, we can define infinite order dominance as follows:

$$\leq_{A_P^\infty} = \bigcup_{r \in \mathbb{N}} \leq_{A_P^r},$$

where  $\mathbb{N} = \{0, 1, \dots\}$ . This means that,

$$y' \leq_{A_P^\infty} y'' \Leftrightarrow y' \leq_{A_P^r} y'' \quad (\text{for some } r \in \mathbb{N}).$$

**Definition 8.** The outcome vector  $y$  is  $A_P^\infty$ -nondominated if and only if there does not exist another outcome vector  $y'$  such that  $y' \leq_{A_P^\infty} y$ . Analogously, a feasible solution  $x$  is called an  $A_P^\infty$ -efficient solution of the problem (2.1) if and only if  $y = f(x)$  is  $A_P^\infty$ -nondominated.

By using Definition 8, we can conclude that  $Y_{A_P^\infty N} = \bigcap_{r \in \mathbb{N}} Y_{A_P^r N}$ .

**Corollary 6.** If  $x \in X$  is an  $A_P^\infty$ -efficient solution for problem (2.1) and  $A_P$  is a matrix with the condition  $a_j \geq 0$  for  $j = 1, \dots, m$ . Then,  $x$  is an efficient solution for problem (2.1).

*Proof.* By applying the definition of  $A_P^\infty$ -efficiency and Corollary 5, the proof is trivial.  $\square$

Corollary 6 indicates that to reduce Pareto optimal solutions, we can use  $A_P^\infty$ -efficient solutions.

## 5. The reduction of $A_P$ -efficient solutions

In this section, our focus is on reducing the set of  $A_P$ -efficient by creating a new preference matrix. This is done by modifying the decomposition of the MOP. As in the previous sections, assume that the partition  $P$  and the preferences matrices  $A_1^P, \dots, A_n^P$  are provided by the DM and  $A_P = A_1^P \oplus \dots \oplus A_n^P$ .

**Definition 9.** Let  $P = \{P_1, \dots, P_n\}$  be a partition of  $\{1, \dots, m\}$  and  $F = \{F_1, \dots, F_t\}$  be a partition of  $\{1, \dots, n\}$ . The generated partition  $E = \{E_1, \dots, E_t\}$  by  $P$  and  $F$  of  $\{1, \dots, m\}$  is defined by  $E_k = \bigcup_{j \in F_k} P_j$  ( $k = 1, \dots, t$ ), where  $F_k$  is index set classes in partition  $P$  should be integrated for class  $k$  in partition  $E$ .

Let  $A_P = A_1^P \oplus \dots \oplus A_n^P$ . If  $F_k = \{k_1, k_2, \dots, k_{|F_k|}\}$  in Definition 9, we define

$$A_k^E = [A_{k_1}^{\prime P} \ A_{k_2}^{\prime P} \ \dots \ A_{k_{|F_k|}}^{\prime P}],$$

where

$$A_{k_j}^{\prime P} = \begin{bmatrix} A_{k_j}^P \\ 0 \end{bmatrix}_{|E_k| \times |P_{k_j}|},$$

for  $j = 1, \dots, |F_k|$ .

**Example 4.** Suppose that  $P_1 = \{1, 2\}$ ,  $P_2 = \{3\}$ ,  $P_3 = \{4\}$ ,  $F_1 = \{1, 2\}$  and  $F_2 = \{3\}$ . The generated partition by  $P$  and  $F$  is  $E_1 = P_1 \cup P_2 = \{1, 2, 3\}$  and  $E_2 = P_3 = \{4\}$ . For example, if  $A_1^P = \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}$ ,  $A_2^P = [1]$  and  $A_3^P = [4]$  then  $A_1^{\prime P} = \begin{bmatrix} 2 & 1 \\ 1 & 3 \\ 0 & 0 \end{bmatrix}$ ,  $A_2^{\prime P} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$  and  $A_1^E = [A_1^{\prime P} \ A_2^{\prime P}] = \begin{bmatrix} 2 & 1 & 1 \\ 1 & 3 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ . Also we have  $A_2^E = A_3^{\prime P} = [4]$ .

In the following, we will investigate the relationship between preference relations  $\leq_{A_P}$  and  $\leq_{A_E}$ .

**Theorem 5.** Let  $P, F, E$  and  $A_P, A_E$  be as in Definition 9. For any two outcome vectors  $y', y'' \in Y$ , we have

$$y' <_{A_P} y'' \Rightarrow y' <_{A_E} y''.$$

Hence  $Y_{A_E N} \subset Y_{A_P N}$ . In particular, if  $P = \{P_1, \dots, P_n\}$  is an arbitrary partition of  $\{1, \dots, m\}$ , then  $Y_{AN} \subset Y_{A_P N}$ , where  $A = [A_1^{\prime P} \ \dots \ A_n^{\prime P}]$  and

$$A_j^{\prime P} = \begin{bmatrix} A_j^P \\ 0 \end{bmatrix}_{m \times |P_j|}.$$

*Proof.* Let  $y' <_{A_P} y''$ . Using Corollary 1, we deduce that

$$A_k^P(y'_{P_k}) \leq A_k^P(y''_{P_k}) \quad (k = 1, \dots, n),$$

and  $A_k^P(y'_{P_k}) \leq A_k^P(y''_{P_k})$  for some  $k \in \{1, \dots, n\}$ . Since  $E_k = \bigcup_{j \in F_k} P_j$ , for all  $k = 1, \dots, t$  and  $A_k^E = [A_{k_1}^{\prime P} \ \dots \ A_{k_{|F_k|}}^{\prime P}]$ , we obtain  $y' <_{A_E} y''$ . Hence  $Y_{A_E N} \subset Y_{A_P N}$ . In particular, let  $P = \{P_1, \dots, P_n\}$  be an arbitrary partition of  $\{1, \dots, m\}$ . For  $F_1 = \{1, \dots, n\}$ , the generated partition by  $P$  and  $F = \{F_1\}$  equal to  $E = \{E_1\}$ , where  $E_1 = \bigcup_{j \in F_1} P_j = \{1, \dots, m\}$ . Now by applying the first part of the theorem, we obtain  $Y_{AN} \subset Y_{A_P N}$ .  $\square$

As shown in the example below, the converse of Theorem 5 is not always true.

**Example 5.** Let  $P_1 = \{1\}$  and  $P_2 = \{2\}$  and  $F_1 = \{1, 2\}$ . The generated partition by  $P$  and  $F$  is  $E_1 = P_1 \cup P_2 = \{1, 2\}$ . If  $A_1^P = A_2^P = [1]$ , then  $A_1^E = [A_1^{P_1} \ A_2^{P_2}] = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ . For  $y' = (2, 1)$  and  $y'' = (1, 3)$ , we have  $y' <_{A_E} y''$ , while  $y' \not\prec_{A_P} y''$ .

In the following, we obtain the relationship between  $A_E$ -efficient solutions and  $A_P$ -efficient solutions by Theorem 5.

**Corollary 7.** *Let  $P, F$  and  $E$  be as in Definition 9 and let  $x \in X$  be a feasible solution. If  $x$  is an  $A_E$ -efficient solution of problem (2.1), then it is an  $A_P$ -efficient solution of problem (2.1). Hence  $X_{A_E E} \subseteq X_{A_P E}$ . Therefore, if  $x \in X$  is an efficient solution of the problem*

$$\begin{aligned} & \min A_E(f(x)) \\ & \text{subject to } x \in X, \end{aligned} \tag{5.1}$$

then it is an efficient solution of the problem (3.4).

According to Theorem 5 and Corollary 7, an algorithm is offered to generate  $A_E$ -efficient solutions, whereby is reduced  $A_E$ -efficient solutions.

---

**Algorithm 1.**

---

Step 1: Determine a partition  $P = \{P_1, \dots, P_n\}$  of  $\{1, \dots, m\}$  and the matrix  $A_P$ , according to the DM.

Step 2: put  $t = 1$ .

Step 3: Consider the partition  $F$ , where  $F_1 = \{1, \dots, t\}$ ,  $F_2 = \{t + 1\}$ ,  $\dots$ ,  $F_{n-t+1} = \{n\}$ .

Step 4: Calculate the partition  $E$  and the matrix  $A_E$ , where  $E_k = \bigcup_{j \in F_k} P_j$  for  $k = 1, \dots, n - t + 1$ , according to Definition 9.

Step 5: Solve the MOP (5.1).

Step 6: If the DM chooses the desired solution, stop.

Step 7: Otherwise put  $t = t + 1$ , if  $t > n$  stop, the model does not answer.

Step 8: Otherwise, go to Step 3.

---

In the first iteration of Algorithm 1, we have  $t = 1$ . Thus  $F_i = \{i\}$  for  $i = 1, \dots, n$ , and hence  $E_k = P_k$  for  $k = 1, \dots, n$ . Therefore, the output of the first iteration of the algorithm is the generation of  $A_P$ -efficient solutions. In particular, if  $P_j = \{j\}$  and  $A_j^P = I$  for  $j = 1, \dots, m$ , then Pareto optimal solutions are computed in the first iteration of this algorithm. For  $t = 2$ , we get  $F_1 = \{1, 2\}$  and  $F_i = \{i + 1\}$  for  $i = 2, \dots, n - 1$ , and hence  $E_1 = P_1 \cup P_2$  and  $E_k = P_{k+1}$  for  $k = 2, \dots, n - 1$ . Hence, in the second iteration of Algorithm 1, the  $A_E$ -efficient solutions are computed. Applying Corollary 7 for these two iterations, we conclude that  $X_{A_E E} \subseteq X_{A_P E}$ . By continuing this process, we observe that the set of solutions obtained in each iteration is reduced compared to the previous iteration, when  $t$  increases to a maximum of  $n$ . In the following example, we obtain the  $A_P$ -efficient solutions where the partition  $P$  of  $\{1, \dots, m\}$  and the matrix  $A_P$  are given by the DM. Then, we gradually reduce




Figure 2. The Pareto optimal and  $A_P$ -efficient solutions with  $P_1 = \{1, 2\}$ ,  $P_2 = \{3, 4\}$ ,  $P_3 = \{5, 6\}$  and  $A_E$ -efficient solutions generated by Algorithm 1.

these solutions by  $A_E$ -efficient solutions in the next iterations of the Algorithm 1. For this purpose, a large number of random solutions are generated for scalable test function. From this large set of solutions, efficient solutions,  $A_P$ -efficient solutions and  $A_E$ -efficient solutions are calculated.

**Example 6.** Let us consider again the test problem (4.2). In Figure 2 from 3000 random solutions, 1804 solutions (blue points) are efficient. Let  $P_1 = \{1, 2\}$ ,  $P_2 = \{3, 4\}$ ,  $P_3 = \{5, 6\}$  and

$$A_1^P = \begin{bmatrix} 1 & 0.5 \\ 0.9 & 0.8 \end{bmatrix}, \quad A_2^P = \begin{bmatrix} 1 & 0.7 \\ 0 & 0.5 \end{bmatrix}, \quad A_3^P = \begin{bmatrix} 0.5 & 0.6 \\ 0.7 & 0.4 \end{bmatrix},$$

be given by the DM. In the first iteration of the Algorithm 1, 1042 solutions (red circles) are  $A_P$ -efficient. 271 solutions (green square) are  $A_{E1}$ -efficient, which are obtained by assuming  $E_1^1 = P_1 \cup P_2 = \{1, 2, 3, 4\}$  and  $E_2^1 = P_3 = \{5, 6\}$ , in the second iteration of the algorithm. Also in the third iteration of the Algorithm 1, 7 solutions (cyan star) are  $A_{E2}$ -efficient, which are obtained by assuming  $E_1^2 = P_1 \cup P_2 \cup P_3 = \{1, 2, 3, 4, 5, 6\}$ .

## 6. Conclusion

The aim of this paper is introducing the new concept of rational  $A_P$ -efficiency for solving of the MOPs, where the preference matrix  $A_P$  is given by DM. This concept is obtained by rational preference relations on the certain cumulative vector  $A_P(y)$  for  $y \in Y$ . We have investigated both the theoretical and practical aspects of rationally



$A_P$ -efficient solutions. Using the powers of  $A_P$ , the concept of  $A_P^r$ -efficiency is expressed to generate a subset of Pareto optimal solutions for  $r = 1, 2, \dots$ . Also, we proved that the  $A_P^r$ -efficient sets are decreasing with respect to  $r$ , and the intersection of these sets is the  $A_P^\infty$ -efficient set. We investigated the relationship between  $A_P$ -efficient and  $A_E$ -efficient solutions, where  $A_E$  is derived from the preference matrix  $A_P$ . Moreover, two experiments were carried out on randomly generated solutions to better compare the efficient solutions with the  $A_P^r$ -efficient solutions and the  $A_P$ -efficient solutions with the  $A_E$ -efficient solutions. These experiments show that the  $A_P^r$ -efficient set is significantly smaller than the efficient set. Furthermore, the  $A_E$ -efficient set is considerably smaller than the  $A_P$ -efficient set.

**Conflict of Interest:** The authors declare that they have no conflict of interest.

**Data Availability:** Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

## References

- [1] V.P. Berman and G.E. Naumov, *Preference-relation with interval value tradeoffs in criterion space*, Autom. Remote Control **50** (1989), no. 3, 398–410.
- [2] S. Dempe, G. Eichfelder, and J. Fliege, *On the effects of combining objectives in multi-objective optimization*, Math. Methods Oper. Res. **82** (2015), 1–18.
- [3] A. Engau and M.M. Wiecek, *Cone characterizations of approximate solutions in real vector optimization*, J. Optim. Theory Appl. **134** (2007), no. 3, 499–513. <https://doi.org/10.1007/s10957-007-9235-8>.
- [4] M. Farina and P. Amato, *On the optimal solution definition for many-criteria optimization problems*, 2002 annual meeting of the North American fuzzy information processing society proceedings. NAFIPS-FLINT 2002 (Cat. No. 02TH8622), IEEE, 2002, pp. 233–238. <https://doi.org/10.1109/NAFIPS.2002.1018061>.
- [5] B.J. Hunt, M.M. Wiecek, and C.S. Hughes, *Relative importance of criteria in multiobjective programming: A cone-based approach*, European J. Oper. Res. **207** (2010), no. 2, 936–945. <https://doi.org/10.1016/j.ejor.2010.06.008>.
- [6] M.M. Kostreva and W. Ogryczak, *Linear optimization with multiple equitable criteria*, RAIRO Oper. Res. **33** (1999), no. 3, 275–297.
- [7] A. Mahmudinejad and D. Foroutannia, *Generalized rationally efficiency in multiobjective programming*, UPB Sci. Bull. A: Appl. Math. Phys. **78** (2016), no. 1, 135–146.
- [8] B. Malakooti, *A decision support system and a heuristic interactive approach for solving discrete multiple criteria problems*, IEEE Trans. Syst. Man. Cybern. **18**

- (1988), no. 2, 273–284.  
<https://doi.org/10.1109/21.3466>.
- [9] J. Molina, L.V. Santana, A.G. Hernández-Díaz, C.A.C. Coello, and R. Caballero, *g-dominance: Reference point based dominance for multiobjective metaheuristics*, European J. Oper. Res. **197** (2009), no. 2, 685–692.  
<https://doi.org/10.1016/j.ejor.2008.07.015>.
- [10] V.D. Noghin, *Relative importance of criteria: a quantitative approach*, J. Multi-Criteria Decis. Anal. **6** (1997), no. 6, 355–363.
- [11] ———, *What is the relative importance of criteria and how to use it in MCDM*, Multiple Criteria Decision Making in the New Millennium: Proceedings of the Fifteenth International Conference on Multiple Criteria Decision Making (MCDM) Ankara, Turkey, July 10–14, 2000, Springer, 2001, pp. 59–68.  
[https://doi.org/10.1007/978-3-642-56680-6\\_5](https://doi.org/10.1007/978-3-642-56680-6_5).
- [12] S. Petchrompo, D.W. Coit, A. Brintrup, A. Wannakrairot, and A.K. Parlikad, *A review of Pareto pruning methods for multi-objective optimization*, Comput. Ind. Eng. **167** (2022), Article ID: 108022.  
<https://doi.org/10.1016/j.cie.2022.108022>.
- [13] S. Petchrompo, A. Wannakrairot, and A.K. Parlikad, *Pruning Pareto optimal solutions for multi-objective portfolio asset management*, European J. Oper. Res. **297** (2022), no. 1, 203–220.  
<https://doi.org/10.1016/j.ejor.2021.04.053>.
- [14] V.V. Podinovskii, *Quantitative importance of criteria*, Autom. Remote Control **61** (2000), 817–828.
- [15] R.E. Steuer, *Multiple Criteria Optimization: Theory, Computation, and Application*, Wiley, New York, 1986.
- [16] V. Venkat, S.H. Jacobson, and J.A. Stori, *A post-optimality analysis algorithm for multi-objective optimization*, Comput. Optim. Appl. **28** (2004), no. 3, 357–372.  
<https://doi.org/10.1023/B:COAP.0000033968.55439.8b>.
- [17] J.B. Yang, *Multiple criteria decision analysis methods and applications*, Hunan Publishing House, Changsha, PR China, 1996, 1996.
- [18] ———, *Minimax reference point approach and its application for multiobjective optimisation*, European J. Oper. Res. **126** (2000), no. 3, 541–556.  
[https://doi.org/10.1016/S0377-2217\(99\)00309-4](https://doi.org/10.1016/S0377-2217(99)00309-4).
- [19] E. Zio, P. Baraldi, and N. Pedroni, *Optimal power system generation scheduling by multi-objective genetic algorithms with preferences*, Reliab. Eng. Syst. Saf. **94** (2009), no. 2, 432–444.  
<https://doi.org/10.1016/j.ress.2008.04.004>.
- [20] E. Zio and R. Bazzo, *A clustering procedure for reducing the number of representative solutions in the Pareto front of multiobjective optimization problems*, European J. Oper. Res. **210** (2011), no. 3, 624–634.  
<https://doi.org/10.1016/j.ejor.2010.10.021>.